



Yield and Characteristics of Melon Fruits under Different Fertilization Management and Soil Cover

**Jackson Teixeira Lobo^{1*}, Witalo da Silva Sales¹, José Fabrício Alves Feitosa²,
Juscelino Martins Costa Junior³, Reynaldo Teodoro de Fatima¹,
Daniel de Almeida Carreiro⁴ and Felipe Thomaz da Camara²**

¹*Program of Postgraduate in Agronomy, Federal University of Paraíba (UFPB), Areia, Brazil.*

²*Federal University of Cariri (UFCA), Crato, Brazil.*

³*Program of Postgraduate in Agroecology and Rural Development, Federal University of São Carlos (UFSCAR), São Carlos, Brazil.*

⁴*Program of Postgraduate in Agronomy, Federal University of São Francisco Valley (UNIVASF), Petrolina, Brazil.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors JTL and WSS designed the study. Authors JTL, WSS, JFAF and JMCJ managed the analyses of the study. Author FTC performed the statistical analysis. Authors JTL and RTF wrote the manuscript. Author DAC managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/v34i530184

Editor(s):

(1) Dr. Ivan Francisco Garcia-Tejero, Instituto Andaluz de Investigacion y Formacion Agraria y Pesquera (IFAPA), Centro Las Torres-Tomejil, Alcala del Rio (Sevilla), Spain.

Reviewers:

(1) Hardarshan Kaur, CMK National P.G. Girls College, Sirsa, India.

(2) Mohamad Awad, National Council for Scientific Research- Centre for Remote Sensing, Lebanon.

(3) Eray Tulukcu, University of Selcuk, Turkey.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48342>

Received 20 January 2019

Accepted 10 April 2019

Published 19 April 2019

Original Research Article

ABSTRACT

Aims: To evaluate the 'Canary' and 'Hale's Best Jumbo' melon yield and fruit characteristics as a function of fertilization management and soil cover with mulch.

Study Design: The design was a randomized complete block design in a 2x3x2 factorial scheme, with three replications and seven plants per plot.

Place and Duration of Study: The experiment was conducted from June to September 2015, at the Center of Agrarian and Biodiversity Sciences of the Federal University of Cariri, located in the city of Crato, Cariri region, Brazil.

*Corresponding author: E-mail: jackson_lob@hotmail.com;

Methodology: Two varieties of melon ('Canary' and 'Hale's Best Jumbo') were studied under three fertilization managements (mineral fertilization via soil, mineral fertilization via soil + organic fertilization and mineral fertilization via soil + foliar fertilization) with the presence or absence of soil cover with polyethylene mulch.

Results: The use of mineral + organic fertilization and mineral + leaf fertilization promoted an increase in fruit mass and yield of approximately 3.93 t ha⁻¹ (25.50%) and 4.64 t ha⁻¹ (30%), respectively. Both melon cultivars presented the best responses grown on plastic mulch, with yield increases of 79.66% and 26.16% for 'Canary' and 'Hale's Best Jumbo', respectively. Moreover, the 'Hale's Best Jumbo' cultivar presented higher soluble solids contents than the 'Canary' melon, with an increase of 11.26% (0.76 ° Brix).

Conclusion: The use of soil cover and additional fertilization (organic or foliar) in the cultivation of melon provides an increase in size, mass and productivity, while soil cover increases the soluble solids content in fruits.

Keywords: *Cucumis melo* L.; 'canary' melon; 'Hale's best jumbo' melon; plant nutrition; mulching.

1. INTRODUCTION

The melon (*Cucumis melo* L.) is an Asian plant of belonging to the family Cucurbitaceae. Fruits are widely appreciated and popular around the world [1]. According to data from the Brazilian Agribusiness Foreign Trade Statistics [2], in 2017, 233.6 thousand tons of melons were shipped, totaling more than 162.9 millions of US\$.

Within the most popular cultivars, the 'Canary' melon belongs to the Inodoros group and has Spanish origin. It presents round yellow fruits with thick and whitish inner flesh, having as main characteristics the resistance to transportation conditions and long post-harvest life, which facilitates the commercialization process [3]. However, in recent years the fruits of aromatic types of melon, such as Cantaloupe, have been gaining more attention of producers and consumers, mainly because they have tastier and earlier fruits. The Cantalupensis variety stands out for bearing spherical and reticulated fruits with salmon colored pulp and intense aroma [4]. The fruits of this group, however, present short post-harvest life, which has hampered the expansion of cultivated areas, signaling the need for research to define the best crop management technologies capable of increasing fruit productivity and quality [5].

In modern agriculture, the search of improvements on yields has led to expressive gains in production and profit, however, some inadequate agricultural practices such as over fertilization can result in significant economic, social and environmental damages [6]. Thus, the use of alternative sources of fertilization may reduce the environmental damage and costs with regular fertilizers [7]. Alternative fertilization

practices, such as foliar feeding, improves the availability of nutrients to the plant, especially micronutrients, in periods of greater demand, which favors the production [8,9]. Moreover, in the last decades, organic fertilization has also been used as an economic and environmental alternative, in the partial or total replacement of chemical fertilizers. Besides releasing nutrients necessary for plant growth, organic fertilizers provide benefits to the soil, such as improvements in structure, aeration and moisture storage capacity, regulating the effect in temperature and cation exchange capacity, which, in turn, potentiates crop productivity [10,11].

An optimum option in the cultivation of several vegetables has been the application of mulching practices. Mulching contributes to the improvement of the production system by reducing temperature fluctuation, loss of water by evaporation and reduces soil erosion. Furthermore, it lessens the direct contact of the fruits with the soil, reducing damages to the rind and improving fruit appearance and quality [12]. Studies on the effects of soil cover on melon cultivation have been developed [13,14], and a better understanding of its association with fertilizer sources becomes important for sustainable management of the crop.

In view of the above stated, the aim of this work was to evaluate the 'Canary' and 'Hale's Best Jumbo' melon yield and fruit characteristics as a function of fertilization management and soil cover with mulch.

2. MATERIALS AND METHODS

The experiment was conducted from June to September 2015, at the Agrarian and Biodiversity

Sciences Center of the Federal University of Cariri, located in the city of Crato, Cariri region, Brazil, (7°14'3.4"S, 39°22'7.6"W; 442 masl). The climate of the region is characterized as Tropical with dry Summer, type As [15], with rainfall that ranges from 700 to 1,000 mm year⁻¹ and an average annual temperature of 27°C.

The soil has a sandy-loam texture, classified as Yellow Red Latosol [16], and the chemical attributes of the 0-20 cm depth are described as follows: pH in water (potential of hydrogen) = 6; O.M.(organic matter) = 4.3 g Kg⁻¹; P (phosphorus) = 3 mg dm⁻³; K⁺(potassium) = 1.3 mmol_c dm⁻³; Na⁺(sodium) = 6.6 mmol_c dm⁻³; Ca²⁺(calcium) = 5 mmol_c dm⁻³; Mg²⁺(magnesium) = 6 mmol_c dm⁻³; Al³⁺(aluminum) = N. D.; H+Al³⁺(hydrogen + aluminum) = 16.5 mmol_c dm⁻³; SB (sum of bases) = 18.9 mmol_c dm⁻³; BS(base saturation) = 53%.

The experimental design was the Randomized Complete Block design in a 2x3x2 factorial scheme, and three replications, totaling 36 plots and seven plants per plot. Studied treatments consisted of two varieties of melon ('Canary' and 'Hale's Best Jumbo') under three fertilization managements (mineral fertilization via soil, mineral fertilization via soil + organic fertilization and mineral fertilization via soil + foliar fertilization) with the presence or absence of soil cover with plastic mulch. Each plot corresponded to a raised garden bed with 0.20 m height, 3.50 m long, 1.80 m wide and area of 6.3 m². In each plot, seven plants were distributed linearly in the center of the garden bed, spaced apart by 45 cm. The useful part consisted of the five central plants of each plot.

Sowing was developed in pits of 30 cm of diameter and 25 cm in depth were dug and the fertilizers corresponding to the treatments were incorporated. Three seeds were sown in each planting hole and the thinning was carried out at 15 DAS (days after sowing) in order to maintain only one plant per pit.

Irrigation was performed by a drip-irrigation system, using flexible a tape with drippers spaced at 30 cm with a flow rate of 1.6 L h⁻¹. The average irrigation time was 2 hours per day, and the amount of water applied was calculated based on the evapotranspiration records observed and according to the Kc coefficient of the melon, defined by Braga Sobrinho et al. [17].

The raised garden beds were covered with black polyethylene mulch after the preparation of the

planting pits and before sowing, maintaining a circular opening measuring 15 cm in diameter.

The mineral fertilization was carried out in all plants regardless of the treatment, based on the soil chemical analysis results and crop requirements, as follows: 40 Kg ha⁻¹ of nitrogen (89 kg ha⁻¹ of urea), 120 kg ha⁻¹ of P₂O₅ (667 kg ha⁻¹ of single superphosphate) and 40 kg ha⁻¹ of K₂O (67 kg ha⁻¹ of potassium chloride).

For the treatment with organic fertilization, four liters of tanned cattle manure were applied per planting pit during the preparation of the raised garden beds (ten days before sowing). The manure presented the following chemical characteristics: pH in water = 8.25; organic matter (OM) = 100.82 g Kg⁻¹; phosphorus (P) = 5.06 mg dm⁻³; potassium (K⁺) = 0.716 mg dm⁻³; sodium (Na⁺) = 1.08 cmol_c dm⁻³; calcium (Ca⁺⁺) = 4 cmol_c dm⁻³; magnesium (Mg⁺⁺) = 3.9 cmol_c dm⁻³; aluminum (Al³⁺) = 0,0; hydrogen + aluminum (H+Al³⁺) = 0.49 cmol_c dm⁻³; cation exchange capacity (CEC) = 10.18 cmol_c dm⁻³; sum of bases (SB) = 9.69 cmol_c dm⁻³; base saturation (BS) = 95.10%.

For the treatment with foliar fertilization, the commercial liquid fertilizer NutrichemCompleto[®] was used at 1% (2L ha⁻¹), and it presents the following composition: N=67.5 g L⁻¹, P₂O₅=108 g L⁻¹, K₂O=67.5 g L⁻¹, Mg=8.1 g L⁻¹, B=5.4 g L⁻¹, Cu=2.7 g L⁻¹, Mn=6.7 g L⁻¹, Zn=13.5 g L⁻¹, TOC=81 g L⁻¹. The first application of leaf fertilizer was performed at 35 DAS, and repeated twice more, at 10 days intervals.

The weeding, turning of the fruits, pest and disease control were carried out at the experimental area according to the recommendations of Braga Sobrinho et al. [17]. Fruit thinning was performed, leaving two fruits per plant. The 'Canary' melon harvesting started at 65 DAS, while the harvest of the 'Hale's Best Jumbo' melon started at 80 DAS. In each variety, five harvests were performed at three days intervals. Fruits were harvested when they presented the formation of the abscission layer of the peduncle [18].

The fruits were weighed on a precision balance, and yield (t ha⁻¹) was estimated based on plant density ha⁻¹. The following variables were measured using a digital caliper: equatorial and polar diameter of the fruits (cm); pulp thickness (cm), corresponding to the average of the thicknesses in the equatorial and polar region

after opening of the fruit to the center in the transverse direction; diameter of the internal cavity (cm), measured in the equatorial region. Soluble solids ($^{\circ}$ Brix) were determined by direct reading in a portable refractometer (model RT-30ATC) according to the norms of the Adolfo Lutz Institute [19].

Data were subjected to analysis of variance to evaluate the effects of the different treatments, applying a Tukey test at 5% probability, according to Banzatto and Kronka [20] recommendation, through the statistical program SISVAR, version 5.3 [21].

3. RESULTS

None of the factors studied in the present research influenced the variable equatorial diameter of the fruits. However, there was an effect of all factors on the polar diameter, in addition to significant interaction between variety and soil cover (mulch) for the same variable. The pulp thickness was affected by the soil cover and fertilization, and the internal cavity diameter affected by the variety and soil cover (Table 1).

The polar diameter of the fruits was higher for the treatments with mineral + organic fertilization and mineral + foliar fertilization, promoting increments of 2.39 and 1.79 cm, respectively, when compared to the average value registered for mineral fertilization only (Table 1).




Fig. 1. Polar diameter of fruits as a function of variety (Canary and Hale's Best Jumbo) and plastic mulch (with or without)

In the bars, the lowercase letters compare the varieties and the uppercase letters compare the soil cover. Bars with the same letters do not differ by Tukey test at 5% probability

When evaluating the interaction between variety and soil cover for the polar diameter of the fruits (Fig. 1), it can be observed that the soil cover promoted a 21.71% increase in the above mentioned variable for 'Canary' melon, while for 'Hales' Best Jumbo' the presence of mulch did not influence this characteristic. When analyzing the variety factor within the soil cover factor, it is noticed that for both covered and uncovered soils the 'Canary' melon obtained the best results for the polar diameter of the fruit

Table 1. Mean square of the variance analysis for Equatorial Diameter (ED), Polar Diameter (PD), Pulp Thickness (PT) and Internal Cavity Diameter (ICD) of 'canary' and 'Hale's Best Jumbo' melons as a function of fertilization management and plastic mulch


Source Variation	ED (cm)	PD (cm)	PT (cm)	ICD (cm)
Variety (V)	1.54 ^{ns}	151.49**	0.11 ^{ns}	7.76**
Canary	10.40	16.9 a	2.89	4.43 b
Hale's Best Jumbo	10.80	12.8 b	3.00	5.36 a
Mulch Cover (MC)	2.82 ^{ns}	40.68**	2.87**	2.73**
With	10.90	15.9 a	3.23 a	5.17 a
Without	10.30	13.8 b	2.67 b	4.62 b
Fertilization (F)	3.66 ^{ns}	13.31**	0.84**	0.21 ^{ns}
Mineral F.	10.10	13.6 b	2.71 b	4.75
Mineral F. + Organic F.	10.50	15.7 a	3.23 a	5.02
Mineral F. + Leaf F.	11.20	15.1 a	2.90 a	4.93
V x MC	0.37 ^{ns}	12.80**	0.19 ^{ns}	0.74 ^{ns}
V x F	0.73 ^{ns}	3.78 ^{ns}	0.05 ^{ns}	0.03 ^{ns}
MC x F	2.59 ^{ns}	0.90 ^{ns}	0.05 ^{ns}	0.48 ^{ns}
V x MC x F	1.54 ^{ns}	3.89 ^{ns}	0.00 ^{ns}	0.19 ^{ns}
Erro	2.99	1.36	0.07	0.23
CV%	16.30	7.88	9.00	9.86

*Means followed by the same lowercase letter in the column do not differ from each other by the Tukey test at 5% probability. **: significant ($P < 0.01$); *: significant ($P \leq 0.05$); ns: non-significant; CV%: coefficient of variation*

Table 2. Mean square of the variance analysis for Fruit Mass (FM), Yield (YLD) and Soluble Solids (SS) of 'Canary' and 'Hale's Best Jumbo' melons as a function of fertilization management and plastic mulch

Source Variation	FM (kg)	YLD (t ha ⁻¹)	SS (°Brix)
Variety (V)	0.07 ^{ns}	39.1263 ^{ns}	5.17 ^{**}
Canary	0.83	19.39	6.75 b
Hale's Best Jumbo	0.74	17.31	7.51 a
Mulch Cover (MC)	0.90 ^{**}	497.6551 ^{**}	6.89 ^{**}
With	0.94 a	22.07 a	7.57 a
Without	0.62 b	14.63 b	6.69 b
Fertilization (F)	0.13 ^{**}	74.8374 ^{**}	0.44 ^{ns}
Mineral F.	0.66 b	15.49 b	6.96
Mineral F. + Organic F.	0.83 a	19.42 a	7.09
Mineral F. + Leaf F.	0.86 a	20.13 a	7.34
V x MC	0.19 ^{**}	106.0350 ^{**}	2.53 ^{ns}
V x F	0.01 ^{ns}	3.0270 ^{ns}	0.56 ^{ns}
MC x F	0.01 ^{ns}	6.9671 ^{ns}	0.17 ^{ns}
V x MC x F	0.02 ^{ns}	9.0668 ^{ns}	0.07 ^{ns}
Erro	0.02	12.7998	0.57
CV%	19.48	19.48	10.6

Means followed by the same lowercase letter in the column do not differ from each other by the Tukey test at 5% probability. ^{**}: significant ($P < 0.01$); ^{*}: significant ($P \leq 0.05$); ^{ns}: non-significant; CV%: coefficient of variation

**Fig. 2. Fruit mass (A) and yield (B) of melon as a function of variety (Canary and Hale's Best Jumbo) and plastic mulch (with or without)**

In the bars, the lowercase letters compare the varieties and the uppercase letters compare the soil cover. Bars with the same letters do not differ by Tukey test at 5% probability

compared to the 'Hale's Best Jumbo' melon, with increments of 40.15% and 23.58% with and without mulch, respectively (Fig. 1).

The pulp thickness of the fruits grown in polyethylene mulched soil presented an average value higher in 0.56 cm than the mean of the uncovered soil. When analyzing the fertilization factor, the lowest average for the pulp thickness was verified for mineral fertilization alone, whereas the addition of organic fertilization increased in 0.52 cm the pulp thickness and the addition of foliar fertilization increased in 0.19 cm (Table 1).

The diameter of the internal cavity in the 'Canary' cultivar was 17.35% lower than the 'Hale's Best Jumbo', and in the uncovered soil there was a 10.64% reduction in comparison to the plastic mulched fruits (Table 1).

Significant interactions were verified between variety and soil cover in relation to fruit mass and yield, showing that there was an interdependence among these factors. It was also observed an isolated effect of fertilization for the two variables (Table 2).

The application of mineral plus organic fertilization promoted an increase in fruit mass and yield of approximately 25.50%, corresponding to an increase of 3.93 t ha⁻¹, while mineral plus leaf fertilization boosted by approximately 30% the averages for these variables, equivalent to 4.64 t ha⁻¹.

Evaluating the interactions between variety and soil cover for fruit mass and yield (Fig. 2), it can be noted that the two melon varieties obtained the best responses when grown on plastic mulch. The 'Canary' and 'Hale's Best Jumbo' melons presented a 79.66% and 26.16% increase in yield, respectively (Fig. 2).

When evaluating the variety factor within the soil cover factor, it is observed that the 'Canary' melon fruit mass and yield were superior to 'Hale's Best Jumbo' when using plastic mulch, with a 28.57% increase in yield; however, without mulching, the melon varieties showed no differences in the responses (Fig. 2).

The 'Hale's Best Jumbo' melon presented higher soluble solids content than the 'Canary' melon, corresponding to a 11.26% increase (0.76 ° Brix), whereas for the soil cover factor, fruits grown on plastic mulch presented 0.88 ° Brix more than fruits grown on uncovered soil (Table 2).

4. DISCUSSION AND CONCLUSION

From the obtained results for the equatorial and polar diameters, it was observed that there was a predominance of growth in 'Canary' melons, which is a feature of this oval-shaped fruit variety, different from 'Hale's Best Jumbo' melons, that have round-shaped fruits. However, the fruit diameters registered in the present study for the 'Canary' melon were inferior to those observed by Dalastra et al. [22], whose values ranged from 12.18 to 13.03 cm for the equatorial diameter and 18.53 to 18.96 cm for the longitudinal (polar) diameter. On the other hand, the 'Hale's Best Jumbo' fruits presented higher diameters (9.83 cm of PD and 8.52 cm of ED) than those found by Rizzo and Braz [23]. According to Dalastra et al. [24], the larger the fruit, the bigger its internal cavity. In agreement with this assertion, those fruits grown on plastic mulch exhibited the greatest cavity measurements, but in the isolated analysis between the varieties it is noticeable that although 'Canary' melons were bigger in size, the average of its internal cavity measurements was

lower than that of 'Hale's Best Jumbo' melons. In this line, it is worth to remark that, those fruits with large internal cavity are less resistant to handling and transportation and suffer greater displacement of the placenta, which leads to a shorter post-harvest shelf life [25]. In the same vein, it is desirable that the fruits have a high pulp thickness, an important feature that makes them more valued in the market [3]. The use of plastic mulch provided better conditions for the fruit pulp development. According to Pinheiro Neto et al. [26], the fact that the mulch keeps soil water from evaporating, thus making it more available to the plants, favors cell division and expansion, which is evidenced by the increased production variables. Braga et al. [27] studied the influence of organic and plastic mulch on the cultivation of melon in Petrolina, Brazil, and did not identify differences in the pulp thickness of mulched and non-mulched fruits. These authors registered 3.87 cm of pulp thickness for fruits grown on polyethylene mulch and 3.35 cm for fruits grown on bare soil, and both results are superior to the ones verified in the present study.

The significant improvements associated to the treatments application, including organic and foliar plus mineral fertilization are due in part to the higher nutritional supply comparing to the use of mineral fertilization only. Both the tanned cattle manure and the foliar fertilizer have in their composition a greater diversity of nutrients that contribute to the development and fruiting of the melon. According to Mantovani et al. [28], besides favoring the chemical properties of the soil by adding organic matter and nutrients, the manure improves the physical and biological properties of the soil. Values similar to those found in the present study were obtained by Nascimento Neto et al. [1] and Charlo et al. [29] for pulp thickness of 'Canary' (3.15 cm) and cantaloupe melon (3.22 cm), respectively. In this line, the application of mineral plus organic fertilization and mineral plus leaf fertilization was beneficial for the fruit mass and yield. The mineral + organic fertilization promoted a slightly lower result than that verified by the mineral plus leaf fertilization; this difference could be understood taking into account the nutrients availability for the plants in each of the treatments. The bovine manure is conditioned to mineralization to release the nutrients which requires time [30], taking into consideration the short cycle of melon production (80 to 90 days) and the application of the organic source close to the sowing date, possibly there was no full use of the nutritional potential of the organic source by

the crop. In melon plants, the leaf constitutes the main source of photoassimilates for the fruits [31], in this sense, the foliar feeding, on the other hand, can give plants a direct boost of nutrients through their leaves, allowing the correction of deficiencies in less time than required by soil fertilization [32].

The interdependence between variety and soil cover for the production variables showed the positive influence of the mulch for the melon crop. Due to inherent features of the studied varieties, there is a tendency for the 'Canary' melon fruits to have higher masses than the 'Hale's Best Jumbo' melons, but from the data in Fig. 2 it is possible to observe that the absence of mulch limited the production, since the two varieties showed no differences. This fact is due to the several benefits of mulching, such as moisture retention, which facilitates nutrient transportation and absorption through the soil solution, weed growth suppression, that keeps unwanted plants from competing with crops for space, light, water and nutrients, as well as the reduction of soil temperature oscillation [12].

When evaluating the influence of mulching on the cultivation of 'Canary' melon, Dantas et al. [14] verified a 145.62% yield increase when the melons were cultivated on plastic mulch compared to cultivation in uncovered soil. Similarly, Morais et al. [33] registered an increase of 82% in yield for mulched 'Canary' melons, a result that is close to those verified in the present study, evidencing the beneficial influence of plastic mulch on the melon cultivation.

Under plastic mulch (Fig. 2A), the mean fruit mass value verified for 'Canary' melon is similar to that found by Dalastra et al. [24] (1.08 kg), while for the average fruit mass of 'Hale's Best Jumbo', a similar value was verified by Vargas et al. [34] (0.84 kg) in Cantaloupe melon. The fruit mass results obtained for both varieties studied in the present research are close to 1 kg, an ideal weigh for marketing in the international trade [29].

When analyzing the results of soluble solids for both varieties, independent of the other evaluated factors, it is observed that the means obtained are lower than those required by the main buyers in the European market (above 9° Brix). The low soluble solids contents verified in the present study are attributed in parts to the application of high water amounts during the

maturation phase of the fruits. According to Pinheiro Neto et al. [26], gradual reduction of the irrigation when approaching the fruit harvest phase is necessary, since a higher water supply causes the dilution of sugars in the plant tissues, leading to a low concentration of soluble solids. During the conduction of the present research, the aforementioned irrigation management was not carried out since the two varieties studied had different production cycles, therefore, different harvest periods, thus the use of a single irrigation system for both of them impeded the reduction of the water volume at the appropriate time for each. When studying the effect of different irrigation levels on 'Gália' melon, Ferraz et al. [35] observed a reduction in soluble solids content with the increase of irrigation volume. Similar results were obtained in the present study, which are in agreement with those reported by Negreiros et al. [13]. These authors registered a 19.6% increase in the soluble solids content of melon fruits grown on polyethylene mulch compared to the fruits grown on bare soil. Moreover, Dalastra et al. [24] recorded higher values of soluble solids content in 'Canary' melon in relation to the 'Cantaloupe' melon, which differs from the results found in this research.

The use of soil cover and additional fertilization (organic or foliar) in the cultivation of melon provides an increase in size, mass and productivity, while soil cover increases the soluble solids content in fruits.

ACKNOWLEDGEMENTS

The authors thank the Federal University of Cariri (UFCA) and the Coordination for the Improvement of Higher Education Personnel (CAPES) for postgraduate scholarships awarding.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nascimento Neto JR, Bomfim GV, Azevedo BM, Viana TVA, Vasconcelos DV. Application methods and nitrogen doses for the yellow melon in Ceará, Brazil. *Irriga*. 2012;17(3):364-375.
2. AGROSTAT/Mapa. Estatísticas de Comércio Exterior do Agronegócio

- Brasileiro/Ministério da Agricultura, Pecuária e Abastecimento. 2018. Exportações Brasileiras 2017. Accessed 20 February 2019. Available: <http://sistemasweb.agricultura.gov.br/pages/AGROSTAT.html>
3. Carmo ILGS, Ferreira RS, Souza JTA, Figueredo LF, Medeiros RD. Production and quality of melon cultivars in Savana de Boa Vista, Roraima. *Rev Agropec Tec.* 2017;38(2):78-83. Available: <http://dx.doi.org/10.25066/agrotec.v38i2.28212>
 4. Zebalos CHS, Soares ER, Barbosa CL, Nogueira AE, Queiroz SF. Liming and fertilization when growing melon. *FAEMA.* 2017;8(2):91-102. Available: <http://dx.doi.org/10.31072/rcf.v8i2.587>
 5. Silva MC, Silva TJA, Bonfim-Silva EM, Farias, LN. Productivity and quality characteristics of nitrogen and potassium fertilized net melon. *Rev Bras Eng Agr Amb.* 2014;18(6):581-587. Available: <http://dx.doi.org/10.1590/S1415-43662014000600003>
 6. Castellanos MT, Cartagena MC, Ribas F, Cabello MJ, Arce A, Tarquis AM. Efficiency indexes for melon crop optimization. *Agron J.* 2010;102(2):716-722. Available: <http://dx.doi.org/10.2134/agronj2009.0286>
 7. Santos APG, Viana TVA, Sousa GG, Gomes-do-Ó LM, Azevedo BM, Santos AM. Yield and quality of melon fruit depending on types and doses of biofertilizers. *Hortic Bras.* 2014;32(4):409-16. Available: <http://dx.doi.org/10.1590/S0102-05362014000400007>
 8. Fernández V, Brown PH. From plant surface to plant metabolism: the uncertain fate of foliar applied nutrients. *Front Plant Sci.* 2013;4:289. Available: <https://doi.org/10.3389/fpls.2013.00289>
 9. Davarpanah S, Tehranifar A, Davarynejad G, Abadfa J, Khorasani, R. Effects of foliar applications of zinc and boron nanofertilizers on pomegranate (*Punica granatum* cv. Ardestani) fruit yield and quality. *Sci Hortic.* 2016;210(10):57-64. Available: <https://doi.org/10.1016/j.scienta.2016.07.003>
 10. Agegnehu G, Nelson PN, Bird MI. Crop yield, plant nutrient uptake and soil physicochemical properties under organic soil amendments and nitrogen fertilization on Nitisols. *Soil Tillage Res.* 2016;160:1-13. Available: <https://doi.org/10.1016/j.still.2016.02.003>
 11. Irineu THS, Figueredo LF, Figueredo JP, Silva JN, Paiva, JRG, Andrade R. Agronomic efficiency of 'Cantaloupe' melon under different water blades and organic fertilization. *Com Sci.* 2018;9(3):421-29. Available: <https://doi.org/10.14295/cs.v9i3.1389>
 12. Braga MB, Resende GM, Moura MSB, Dias RCS, Costa ND, Calgaro M, Correia JS, Silva FZ. Productivity and quality under different types of cover soil. *Irriga.* 2010;15(4):422-30.
 13. Negreiros MZ, Costa FA, Medeiros JF, Leitão MVBR, Bezerra Neto F, Espínola Sobrinho J. Yield and fruit quality of melon cultivated under irrigation depths and coverage with polyethylene film of different colors. *Hortic Bras.* 2005;23(3):773-779. Available: <https://doi.org/10.1590/S0102-05362005000300017>
 14. Dantas DC, Medeiros JF, Freire AG. Yield and quality of the melon fruit grown with plastic films in response to irrigation depth. *Cienc Agron.* 2011;42(3):652-661.
 15. Alvares, CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's climate classification map for Brazil. *Meteorol Z.* 2013;22(6):711-728.
 16. FUNCEME (Fundação Cearense de Meteorologia e Recursos Hídricos). Levantamento de reconhecimento de média intensidade de solos: mesorregião do sul cearense. Fortaleza, 2012. 280p. Accessed 25 February 2019. Available: <http://www.funceme.br/areas/17-mapas-tem%C3%A1ticos/544-solos/>.
 17. Braga Sobrinho R, Assis JS, Guimarães JA, Freitas JAD, Bastos MSR. Produção integrada de melão. Fortaleza: Embrapa Agroindústria Tropical; 2008.
 18. Queiroga RCF, Puiatti M, Fontes PCR, Cecon PR. Yield and quality of muskmelon fruits varying fruit and leaf numbers per plant. *Hortic Bras.* 2008;26(2):209-215. Available: <http://dx.doi.org/10.1590/S0102-05362008000200016>
 19. Instituto Adolfo Lutz. Métodos Físico-químicos para análise de alimentos. São Paulo: Instituto Adolfo Lutz; 2008.
 20. Banzatto DA, Kronka SN. Experimentação agrícola. 1st ed. Jaboticabal: FUNEP; 1989.

21. Ferreira DF. Sisvar: A computer statistical analysis system. *Cienc Agrotec.* 2011; 35(6):1039-42.
Available:<http://dx.doi.org/10.1590/S1413-70542011000600001>
22. Dalastra GM, Echer MM, Hachmann TL. Performance of cultivars of melon, depending on the number of fruits per plant. *JAS.* 2015;4(1):26-41.
23. Rizzo AAN, Braz LT. Evaluation of qualitative characteristics of netted melon fruits under greenhouse conditions. *Hortic Bras.* 2001;19(3):237-240.
Available:<http://dx.doi.org/10.1590/S0102-05362001000300017>
24. Dalastra GM, Echer MM, Klosowski ES, Hachmann TL. Yield and quality of three types of melon, varying the number of fruits per plant. *Ceres.* 2016;63(4):523-531.
Available:<http://dx.doi.org/10.1590/0034-737X201663040013>
25. Dantas DJ, Mendonça V, Nunes, GHS, Guimarães IP, Dantas DJ. Evaluation of yield and quality of cantaloupe melon hybrids. *Rev Verde.* 2011;6(4):132-136.
26. Pinheiro Neto LG, Viana TVA, Azevedo BM, Freitas JAD, Souza, VF. Melon fruit production and quality under hydric reduction. *Irriga.* 2007;12(1):54-62.
Available:<http://dx.doi.org/10.15809/irriga.2007v012n1p54-62>
27. Braga MB, Marouelli WA, Resende GM, Moura MSB, Costa ND, Calgaro M, Correia JS. Soil covers on the melon cultivation under nonwoven blanket. *Hortic Bras.* 2017;35(1):147-153.
Available:<http://dx.doi.org/10.1590/S0102-053620170123>
28. Mantovani JR, Silveira LG, Landgraf PRC, Santos AR, Costa BS. Phosphorus rates and use of cattle manure in potted gerbera cultivation. *Ornam Hortic.* 2017;23(4):412-418.
Available:<https://doi.org/10.14295/oh.v23i4.1012>
29. Charlo HCO, Galatti FS, Braz LT, Barbosa JC. Net melon experimental hybrids cultivated in soil and substrate. *Rev Bras Frutic.* 2011;33(1):144-156.
Available:<http://dx.doi.org/10.1590/S0100-29452011005000028>
30. Souto AGL, Cavalcante LF, Silva MRM, Ferreira Filho RM, Lima Neto AJ, Diniz BLMT. Nutritional status and production of noni plants fertilized with manure and potassium. *J Soil Sci Plant Nutr.* 2018; 18(2):403-417.
Available:<http://dx.doi.org/10.4067/S0718-95162018005001301>
31. Castellanos MT, Cabello MJ, Cartagena MC, Tarquis AM, Arce A, Ribas F. Growth dynamics and yield of melon as influenced by nitrogen fertilizer. *Sci Agr.* 2011;68(2): 191-99.
Available: <http://dx.doi.org/10.1590/S0103-90162011000200009>
32. Fageria NK, Barbosa Filho MP, Moreira A, Guimarães CM. Foliar Fertilization of Crop Plants. *J Plant Nutr.* 2009;32(6):1044-64.
Available:<https://doi.org/10.1080/01904160902872826>
33. Morais ERC, Maia CE, Negreiros, MZ, Araújo Junior, BB, Medeiros, JF. Growth and yield of the Goldex Melon crop influenced by coverage of the soil. *Sci Agrar.* 2008;09(2):129-137.
Available:<http://dx.doi.org/10.5380/rsa.v9i2.10940>
34. Vargas PF, Galatti FS, Souza JO, Castoldi R, Charlo HCO, Braz LT. Physicochemical characteristics of experimental net melon hybrids developed in Brazil. *Hortic Bras.* 2013;31(3):351-355.
Available: <http://dx.doi.org/10.1590/S0102-05362013000300002>
35. Ferraz RLS, Melo AS, Ferreira RS, Dutra AF, Figueredo LF. Morpho-physiological aspects, yield and water-use efficiency in the Gália melon plant in a protected environment. *Cienc Agron.* 2011;42(4):957-964.
Available:<http://dx.doi.org/10.1590/S1806-66902011000400018>

© 2019 Lobo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48342>