



# Effect of Boron, Zinc and Sulphur on Growth, Yield and Mineral Composition of Black Gram (*Vigna mungo* L.) in Alluvial Soil

Kavita, Y.V. Singh <sup>a\*</sup>, S.K. Singh <sup>b</sup>, P.K. Bharteey <sup>c</sup>,  
Kajal Singh <sup>a</sup>, Ashutosh Sahu <sup>a</sup> and Aman Saini <sup>a</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi-221011, Uttar Pradesh, India.

<sup>b</sup> Department of Soil Science, Tirhut College of Agriculture, Dholi, Dr. RPCAU, PUSA, Samastipur- 848125, Bihar, India.

<sup>c</sup> Department of Agricultural Chemistry and Soil Science, C.C.R. (P.G.) College, Muzaffarnagar-251 001, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i84884>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120712>

Original Research Article

Received: 25/05/2024

Accepted: 01/08/2024

Published: 05/08/2024

## ABSTRACT

The present study determining the effect of Boron, Zinc and Sulphur on Growth, Yield and Mineral Composition of Black gram (*Vigna mungo* L.) in alluvial soil. A pot experiment was conducted during kharif 2023 at Net House in Department of Soil Science and Agricultural Chemistry, Institute

\*Corresponding author: E-mail: [yvsingh59@rediffmail.com](mailto:yvsingh59@rediffmail.com);

of Agricultural Sciences, B.H.U, Varanasi, U.P. to study the effect of Boron, Zinc and Sulphur on growth, yield and quality of black gram (*Vigna mungo* L.). The experiment was laid-out in a Completely Randomized design (CRD) with eight treatments, each having three replications. Dry matter accumulation, Number of branches plant<sup>-1</sup>, test weight, grain yield kg ha<sup>-1</sup>, straw yield kg ha<sup>-1</sup>, harvest index significantly affected due to various treatments. T<sub>7</sub> (B +Zn+ S) recorded most pronounced improvements.

**Keywords:** Blackgram; sulphur; zinc; boron; test weight; quality and yield.

## 1. INTRODUCTION

Pulses are integral to Indian cooking due to their affordability, accessibility, and rich protein content. Farmers benefit significantly from cultivating pulse crops, which offer sustainable agricultural advantages such as minimal water needs, deep root systems that prevent soil erosion, extensive ground coverage, and their ability to enhance productivity and resilience in farming through crop rotation and intercropping [1]. Pulse crops hold a crucial role in Indian agriculture for several reasons. Firstly, they have the unique capability of biological nitrogen fixation, which enriches the soil with essential nutrients. Additionally, their deep root systems help in mobilizing insoluble soil nutrients and improving soil structure. These plants also bring about beneficial changes in the physical properties of the soil, earning them the title of 'soil fertility restorers'. Beyond their agronomic benefits, pulses are highly valued as sources of food, fodder, and animal feed. India is the world's largest producer of pulses, accounting for one-third of the global pulse acreage and a quarter of its total production. For a significant segment of the population, especially vegetarians, pulses are the main source of protein [2,3].

Black gram (*Vigna mungo* L.) is a significant pulse crop and a staple in tropical and subtropical regions. Also known as urad bean, urad dal, or urad, it thrives in light soils that retain moisture, though loamy and clay loam soils are also suitable. Black gram (*Vigna mungo* L.) is a significant pulse crop and a staple in tropical and subtropical regions. Also known as urad bean, urad dal, or urad, it thrives in light soils that retain moisture, though loamy and clay loam soils are also suitable. Neutral pH loam to clay loam soils is ideal for its cultivation, as it does not tolerate waterlogged conditions well. Black gram is valued for its nutritional richness, boasting high protein content (22.24%), carbohydrates (56.6-59.6%), fats (1.2-1.4%), minerals (3.2%), and significant amounts of phosphorous (385 mg/100g), calcium, and iron [1]. Unlike other

pulses, it stands out for its somewhat mucilaginous texture, which adds density due to its long polysaccharide carbohydrate chains. It provides affordable protein, it is often referred to as the "poor man's meat" [4,5,6].

Boron is an essential element for healthy plant growth and development. It is essential to the fertilization and blooming processes, which raise crop yield and quality [7,8,9]. Since its deficiency has been documented in 132 crops, it is acknowledged as one of the most often deficient micronutrients in soils. Zinc is a crucial nutrient for plants that is involved in several metabolic, regulatory, and developmental processes [7]. Additionally, it has been demonstrated to play a role in defense against the harmful effects of reactive oxygen species [10,11]. Concern over zinc nutrition in plants is growing as zinc deficiency is common in the world's arid and semi-arid regions [12]. Sulphur (S) is a vital plant nutrient necessary for the manufacture of protein, vitamins, enzymes, and amino acids such as cystine, cysteine, and methionine. It also plays a major role in maintaining increased output of pulse crops and participates in several metabolic and enzymatic processes, including respiration, photosynthesis, and the nitrogen fixation symbiotic relationship between legumes and rhizobium [13].

## 2. MATERIALS AND METHODS

### 2.1 Experimental Site

A pot experiment on black gram was conducted on July 30<sup>th</sup>, 2023 in the Net House of the Department of Soil Science and Agricultural Chemistry at the Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, situated at an altitude of 80.71 meters above mean sea level, lies between 25°18' North latitude and 80°36' East longitude. "The region has a humid subtropical climate, with an average annual rainfall of approximately 1100 mm and a potential evapotranspiration of around 1525 mm" [14].

**Table 1. Treatment details of the experiment in black gram**

Treatment details	Abbreviations
Control	T <sub>0</sub>
RDF+Boron @ 1.5 kg ha <sup>-1</sup>	T <sub>1</sub>
RDF+Zinc @ 5 kg ha <sup>-1</sup>	T <sub>2</sub>
RDF+Sulphur @ 20 kg ha <sup>-1</sup>	T <sub>3</sub>
RDF+Boron@1.5 kg ha <sup>-1</sup> + Zinc(@5 kg ha <sup>-1</sup>	T <sub>4</sub>
RDF+Boron@ 1.5 kg ha <sup>-1</sup> + Sulphur(@ 20 kg ha <sup>-1</sup>	T <sub>5</sub>
RDF+Zinc@ 5 kg ha <sup>-1</sup> + Sulphur(@ 20 kg ha <sup>-1</sup>	T <sub>6</sub>
RDF+Boron@1.5 kg ha <sup>-1</sup> +Zinc @ 5 kg ha <sup>-1</sup> +Sulphur(@ 20 kg ha <sup>-1</sup>	T <sub>7</sub>

Where: Recommended dose of fertilizer N, P and K, seed treatment with Rhizobium + PSB was given for each treatment, Recommended dose of fertilizer N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O, 20: 40: 20 kg ha<sup>-1</sup>[15]

## 2.2 Experimental Setup

Initially, the soil had a sandy clay loam texture. It was slightly alkaline (pH 7.41), had normal salt content (EC 0.17 dS m<sup>-1</sup>), low organic carbon content (0.48%), low available nitrogen (155.36 kg ha<sup>-1</sup>), high available phosphorus (23.46 kg ha<sup>-1</sup>), and medium available potassium (190 kg ha<sup>-1</sup>) and Available Sulphur (17.80 kg ha<sup>-1</sup>). The soil also had DTPA-extractable micronutrients with concentrations of Zn at 1.20 mg kg<sup>-1</sup> and Available boron 0.51 mg kg<sup>-1</sup>. During August-September 2023, a pot experiment was conducted using Black gram (variety T-9) as the test crop. The soil for the experiment was collected from the Agro farm at BHU, air-dried, ground, sieved through a 2-mm mesh, and placed in polythene-lined pots. The experiment was laid-out in a Completely Randomized design (CRD) with eight treatments, each having three replications. Uniform basal application was made with 20 kg N ha<sup>-1</sup>, 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, 20 kg K<sub>2</sub>O ha<sup>-1</sup> through urea, DAP (Diammonium phosphate) and MOP (Murate of Potash), respectively. Sulphur, Zn and B were applied @ 20, 5 and 1.5 kg ha<sup>-1</sup> from Bentonite-S, Zinc sulphate and Di-sodium tetra-borate pentahydrate, respectively. The pots were kept at field capacity throughout the study period.

## 2.3 Growth and Yield Attributes

Growth attributing characters of blackgram mainly plant height at 20, 40 and 60 days after sowing (DAS), number of branches per plant, dry matter accumulation were recorded. At maturity grain yield, straw yield, test weight (g) was recorded and harvest index were calculated.

$$\text{Harvest index} = \frac{\text{Economical yield}}{\text{Total biological yield}} \times 100$$

## 2.4 Statistical Analysis

"The research data were analyzed using statistical software SPSS 16.0 for ANOVA (complete randomized design). Duncan multiple range test (DMRT) at p≤0.05 levels of significance was used to evaluate the significant differences among mean values" [16].

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters

The analysis of variance for the data on plant height shows non-significant difference was found on growth stages .T<sub>7</sub> (Boron + Zinc + Sulphur) exhibited the highest plant height (17.31,39,45.69 cm) which was statistically at par with T<sub>5</sub> (Boron + Sulphur) that is (17.05,39.11,44.18 cm).This was followed by treatment T<sub>6</sub> (Zinc + Sulphur), T<sub>4</sub> (Boron + Zinc), and T<sub>3</sub> (Sulphur @20 kg ha<sup>-1</sup>), with harvest plant heights of 43.8 cm, 43.34 cm, and 42.58 cm, respectively. In contrast, the control group recorded the lowest harvest plant height of 40.75 cm T<sub>7</sub> was statistically at par with T<sub>5</sub> and T<sub>6</sub>.More availability of B, Zn and S might have encouraged growth promoting hormones and protein synthesis means more chlorophyll content leading to higher photosynthesis. These resulted into rapid cell elongation and enlargement and ultimately plant height [17,18,14]. Number of branches count plant<sup>-1</sup> was statistically significant due to various treatments (Table 2).

The analyzed data relating to the dry matter accumulation at harvest of black gram is shown in Table 1. T<sub>7</sub>, which involved the application of boron, zinc, and sulphur, resulted in the highest dry matter accumulation at 4.95 grams per plant. This was followed closely by treatment T<sub>5</sub>, which involved the application of boron and sulphur,

with 4.78 grams per plant. Treatments T<sub>6</sub> (zinc and sulphur) and T<sub>4</sub> (boron and zinc) recorded 4.56 grams per plant and 4.50 grams per plant, respectively. In contrast, the lowest dry matter accumulation of 3.98 grams per plant was observed in treatment T<sub>2</sub>, which involved zinc application at 5 kg ha<sup>-1</sup>. T<sub>7</sub> was statistically at par with all the treatments except the control. The data revealed a significant difference among the various treatments. These findings align with previous research by [19], which found that the sulphur rate significantly affects soybean dry matter.

The highest of branches plant<sup>-1</sup> was observed with application of B + Zn + S, which was statistically superior from all the treatments except T<sub>5</sub> (B + S). The lowest number of branches plant<sup>-1</sup> was observed with control T<sub>0</sub> [20,15].

### 3.2 Yield Component

The data of grain yield were analyzed and summarized in Table 3. Different treatments exhibited significant variation in grain yield of urad bean. Treatment T<sub>7</sub>, involving the application of boron, zinc, and Sulphur, resulted in the highest seed output at 1.95 t ha<sup>-1</sup> followed by T<sub>5</sub> (1.68 t ha<sup>-1</sup>) and T<sub>6</sub> (1.62 t ha<sup>-1</sup>). In contrast, the control treatment (T<sub>0</sub>) exhibited the lowest grain production at 0.66 t ha<sup>-1</sup>. T<sub>7</sub> demonstrated the highest haulm yield of 3.71 t ha<sup>-1</sup>, surpassing the control (2.61 t ha<sup>-1</sup>) and treatments T<sub>5</sub> (3.63 t ha<sup>-1</sup>), and T<sub>6</sub> (3.5 t ha<sup>-1</sup>). T<sub>7</sub> was statistically at par with T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub>. However, three other treatments with application of B + S,

Zn + S and B + Zn were also found effective to enhance yield over control as well as those treatments where applied alone nutrients i.e. B, Zn and S. The lowest grain and straw yield were found in control. The Harvest Index of black gram demonstrates significant variation ranging from 20.21% to 34.40%. T<sub>7</sub>, exhibited the highest Harvest Index at 34.45 %, while the control treatment (T<sub>0</sub>) showed the lowest Harvest Index at 20.21% [17]. Treatments exerted significant impacts on variations in test weight (1000-seed weight). T<sub>7</sub> (Boron + Zinc + Sulphur) emerged as the most effective, producing the highest test weight of 40.20 grams followed by T<sub>5</sub> and T<sub>6</sub>. Conversely, the control group (T<sub>0</sub>) exhibited the lowest test weight at 33.20 grams. Noteworthy improvements in test weight were observed with alternative treatments.

### 3.3 Nutrient Content

Table 4 shows Nitrogen content ranged from 1.80% to 3.80% in seeds and 1.40% to 2.80% in haulm across the treatments. Analysis of the data revealed that nitrogen content increased in both seed and haulm compared to the control when treated with boron, Zinc and Sulphur. T<sub>0</sub> had the lowest content of phosphorus (0.51 and 0.21), in seed and haulm while the T<sub>7</sub> with combine application of Boron, Zinc and Sulphur had the highest content of Phosphorus (1.17 and 0.45), followed by the T<sub>5</sub>, T<sub>6</sub> and T<sub>4</sub>. The T<sub>0</sub> produced the lowest value (0.20 % and 0.41%) in seed and haulm, respectively, whereas the combine application of Boron, Zinc and Sulphur (T<sub>7</sub>) had the highest potassium content in both seed and haulm (1.04% and 0.89%).

**Table 2. Effect of different treatments on plant height, dry matter accumulation and number of branches per plant at harvest of black gram**

Treatments	Plant height (cm)			Dry matter accumulation (g plant <sup>-1</sup> )	Number of branches per plant at harvest
	at 20 DAS	at 40DAS	at 40DAS		
Control	13.51	34.01	40.75	3.98	6.23
T <sub>0</sub>	16.10	36.35	42.06	4.35	7.16
T <sub>1</sub>	15.76	36.06	41.85	4.20	6.67
T <sub>2</sub>	16.03	36.65	42.58	4.38	7.42
T <sub>3</sub>	16.10	37.02	43.34	4.50	7.48
T <sub>4</sub>	17.05	39.11	44.18	4.78	7.82
T <sub>5</sub>	16.83	38.78	43.80	4.56	7.64
T <sub>6</sub>	17.31	39.00	45.69	4.95	8.23
T <sub>7</sub>	0.67	1.211	1.89	0.15	0.21
CD (5%)	1.33	1.94	1.80	0.54	0.41

### 3.4 Available Nutrients in Post-Harvest Soil of Black Gram

Table 5 shows the lowest amount of nitrogen, at 160 kilograms per hectare, was observed in the control (T<sub>0</sub>), while the soil in T<sub>7</sub> exhibited the significantly highest nitrogen content at 296.00 kg per hectare. T<sub>7</sub> resulted in the largest amount

of accessible phosphorus (50.90 kg ha<sup>-1</sup>). The available phosphorus value was lowest (23.06 kg ha<sup>-1</sup>) in the control, or T<sub>0</sub>. The data in Table 5 shows information on available potassium in soil post the harvest of Black gram and it varied from 201.15 to 312.20 kg ha<sup>-1</sup> indicating that soil was categorized under medium to high in available potassium.

**Table 3. Effect of different treatments on seed yield, haulm yield, harvest index and test weight of black gram.**

Treatments	Seed yield (t ha <sup>-1</sup> )	Haulm Yield (t ha <sup>-1</sup> )	Harvest index (%)	Test weight (gm)
T <sub>0</sub>	0.66	2.61	20.2	33.20
T <sub>1</sub>	1.03	3.06	25.06	34.20
T <sub>2</sub>	0.82	2.8	22.73	32.80
T <sub>3</sub>	1.35	3.17	29.9	34.70
T <sub>4</sub>	1.56	3.41	31.32	37.20
T <sub>5</sub>	1.68	3.63	32.72	38.75
T <sub>6</sub>	1.62	3.55	31.65	38.21
T <sub>7</sub>	1.95	3.71	34.45	40.20
SEm±	0.03	0.139	0.763	0.78
CD (0.05)	0.1	0.42	4.48	2.24

**Table 4. Effect of different treatments on nutrient contents in seed and haulm of black gram**

Treatments	N Content (%)		P Content (%)		K Content (%)	
	Seed	Haulm	Seed	Haulm	Seed	Haulm
T <sub>0</sub>	1.80	1.40	0.51	0.21	0.20	0.41
T <sub>1</sub>	2.68	1.91	0.81	0.24	0.39	0.57
T <sub>2</sub>	2.15	1.60	0.69	0.23	0.37	0.53
T <sub>3</sub>	2.98	1.94	0.98	0.34	0.42	0.64
T <sub>4</sub>	3.12	1.98	1.05	0.37	0.51	0.69
T <sub>5</sub>	3.80	2.60	1.11	0.41	0.80	0.84
T <sub>6</sub>	3.64	2.27	1.08	0.38	0.74	0.77
T <sub>7</sub>	3.90	2.80	1.17	0.45	1.04	0.89
SEm±	0.059	0.021	0.073	0.005	0.012	0.371
CD (0.05)	0.19	0.05	0.053	0.014	0.042	1.14

**Table 5. Effect of different treatments on available N, P, K, S (kg ha<sup>-1</sup>) and B (mg kg<sup>-1</sup>) in post-harvest soil of black gram**

Treatments	Available Nitrogen (kg ha <sup>-1</sup> )	Available Phosphorus (kg ha <sup>-1</sup> )	Available potassium (kg ha <sup>-1</sup> )	Available Sulphur (kg ha <sup>-1</sup> )
T <sub>0</sub>	167.00	23.06	201.15	21.45
T <sub>1</sub>	225.20	27.89	235.48	23.45
T <sub>2</sub>	215.34	25.64	224.45	25.65
T <sub>3</sub>	260.50	31.15	250.54	30.86
T <sub>4</sub>	277.28	32.46	262.34	29.45
T <sub>5</sub>	283.20	42.67	290.89	31.45
T <sub>6</sub>	281.06	36.78	276.60	32.86
T <sub>7</sub>	296.00	50.90	312.20	35.68
SEM±	2.17	1.04	2.74	0.79
CD 5%	6.54	3.20	8.28	2.45

Where: T<sub>0</sub>=Control, T<sub>1</sub>=RDF+B @1.5 kg ha<sup>-1</sup>, T<sub>2</sub>=RDF+Zn @5kg ha<sup>-1</sup>, T<sub>3</sub>= RDF+S@20 kg ha<sup>-1</sup>, T<sub>4</sub>=RDF+B+Zn, T<sub>5</sub>=RDF+B+S, T<sub>6</sub>=RDF+Zn+S, T<sub>7</sub>=RDF+B+Zn+SRDF=Recommended dose of fertilizer

#### 4. CONCLUSION

Increased plant height, accumulation of dry matter, and more branches were observed upon application of (B@ 1.5 kg ha<sup>-1</sup>, Zn (@ 5 kg ha<sup>-1</sup>, S 20 kg ha<sup>-1</sup>). When T<sub>7</sub>(B+Zn+ S) was applied in combination, the maximum grain yield, straw yield, harvest index, and test weight were also recorded which was significantly superior over T<sub>0</sub> (control), T<sub>1</sub> (B), T<sub>3</sub> (S), T<sub>4</sub> (B + Zn), and T<sub>2</sub> (Zn) and at par with T<sub>5</sub> (B + S) and T<sub>6</sub> (Zn + S). Though all treatments showed positive effect in increasing growth parameter and yield. This suggests that the application of these nutrients enhances the nutritional quality of the seeds, making them more valuable in terms of human and animal nutrition. Additionally, the study highlights improvements in soil properties resulting from the application of boron, zinc and sulphur. The treatment leads to enhancements in soil fertility, indicated by higher levels of organic carbon and nitrogen content in the soil. This suggests that the combination of these nutrients not only benefits the crop but also contributes to long-term soil health and sustainability.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Singh RE, Singh V, Tiwari D, Masih A. Effect of levels of phosphorus and sulphur on growth and yield of black gram (*Vigna mungo* L.). *International Journal of Current Microbiology and Applied Science*. 2020;9 (10):2784-2791.
2. Ghosh K, Swaroop N, Thomas T, Ravindra J. Soil physico-chemical properties as Influenced by combined use of NPK and zinc at varying levels under blackgram (*Vigna mungo* L.) cultivation in an inceptisol of Prayagraj, Uttar Pradesh, India. *International Journal of Plant & Soil Science*. 2022;34(22):1172-1182.
3. Banerjee P, Kumari VV, Nath R. Response of autumn sown black gram as influenced by cobalt, potassium and boron nutrition under Indian subtropics. *Journal of Plant Nutrition*. 2023;46(19):4573-4593.
4. Devi SS, Tuppad P, Kishore A. A review of the effects of phosphorus and sulphur levels on growth, yield, and quality of gram and beans: Phosphorus and sulphur levels on gram and beans. *The Journal of Rural Advancement*. 2022;10(01):19-28.
5. Gowthami P. Effect of foliar application of potassium, boron and zinc on growth and yield of soybean (*Glycine Max* L.) (Merrill) (Doctoral Dissertation, Acharya NG Ranga Agricultural University, Guntur. 2014;4(3): 81-86
6. Sahu S, Singh PK. Effect of micronutrients and biofertilizer inoculation on grain yield, protein content, micronutrients content and economics of chickpea. *Annals of Agricultural Research*. 2009;30:(1-2).
7. Aslam M, Hussain N, Zubair M, Hussain SB, Baloch MS. Integration of organic and inorganic sources of phosphorus for Increased productivity of mung bean (*Vigna radiata* L). *Pakistan Journal of Agriculture Science*. 2010;7(2):111-114.
8. Kanwar JS, Randhawa NS. Micronutrient research in soils and Plants in India: A review. *ICAR. Technical Bulletin (Agriculture)*. 1974;50:1-60.
9. Shorrocks VM. The occurrence and Correction of boron deficiency. *Plant and Soil*. 1997;(193): 121-148.
10. Shi GR, Cai QS. Photosynthetic and anatomic responses of peanut leaves to zinc stress. *Biologia Plantarum*. 2009; (53):391-394.
11. Ram S, Katiyar TP. Effect of sulphur and zinc on the seed yield and protein content of summer mungbean under arid climate. *International Journal of Science and Nature*. 2013;4(3):563-6.
12. Cakmak I. Tansley Review No. 111 Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *The New Phytologist*. 2000; 146(2):185-205.
13. Takkar PN, Walker CD. The distribution and correction of zinc deficiency. In *Zinc in soils and plants: Proceedings of the International Symposium on 'Zinc in Soils and Plants' held at The University of Western Australia*. Dordrecht: Springer Netherlands. 1993;151-165.
14. Kannan P, Arunachlam P, Prabukumar G, Prabhakaran J. Response of blackgram (*Vigna mungo* L.) to multi-micronutrient

- mixtures under rainfed alfisol. Journal of the Indian Society of Soil Science. 2014;62(2):154-160.
15. Patel S. Role of boron and sulfur in enhancing soil fertility and black gram yield. Agronomy Journal, 2021;113(2):567-578.
  16. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & sons; 1984.
  17. Rao CS, Singh KK, Masood Ali MA. Sulphur: A key nutrient for higher pulse production. 2001; 46:37-48.
  18. Kaisher MS, Rahman M.A.S, Amin M.H.A, Amanullah A.S.M, and Ahsanullah A. S. M. Effect of sulphur and boron on the seed yield and protein content of mungbean. Bangladesh Research publication Journal. 2010;3(4):1181-1186.
  19. Ahmad I, Akhtar MJ, Asghar HN, Khalid M. Influence of rhizobium applied in combination with micronutrients on mungbean. Pakistan Journal of Life and Social Sciences. 2013;11(1):53-59.
  20. Najjar GR, Singh SR, Akhtar F, Hakeem SA. Influence of sulphur level on yield, uptake and quality of soybean (Glycine max) under temperate conditions of Kashmir valley. Indian Journal of Agricultural Sciences. 2011;81(4):340.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/120712>