



Anaemia in Pregnancy at Booking: Prevalence and Risk Factors among Antenatal Attendees in a Southern Nigeria General Hospital

Victory E. Iyanam^{1*}, Alphonsus U. Idung¹, Henry E. Jombo²
and Ndifreke E. Udonwa¹

¹Department of Family Medicine, Faculty of Clinical Sciences, University of Uyo, Nigeria.

²Department of Psychiatry, Faculty of Clinical Sciences, University of Uyo, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2019/v15i130112

Editor(s):

(1) Dr. John K. Triantafyllidis, Associate Professor, Iasi University of Medicine and Pharmacy, Romania and IASO General Hospital, Holargos, Athens, Greece.

Reviewers:

(1) Shigeki Matsubara, Jichi Medical University, Japan.
(2) Nsiah Kwabena, Kwame Nkrumah University of Science and Technology, Ghana.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/48805>

Received 19 February 2019

Accepted 28 April 2019

Published 04 May 2019

Original Research Article

ABSTRACT

Background: Despite the fact that anaemia is a preventable morbidity in most cases, its prevalence among pregnant women is still unacceptably high especially in rural and sub-urban settings. This is worrisome considering the enormous contribution of anaemia to maternal and infant morbidity and mortality.

Objective: This study aimed at determining the prevalence of anaemia at booking, identifying and describing its risk factors among pregnant women who attended the antenatal clinic of General Hospital, Etinan, a sub-urban area in Akwa Ibom State, South-South Nigeria.

Methodology: This was a cross-sectional descriptive study involving 375 pregnant women seen at the antenatal (booking) clinic of General Hospital, Etinan, between April and October 2018, recruited through non-probability consecutive sampling technique. A pretested semi-structured interviewer-administered questionnaire was used to obtain data on socio-demographic, family and nutritional characteristics and obstetric and medical history of the respondents. Blood sample of each respondent was obtained from the median cubital vein, analysed with haemoglobinometer and their haemoglobin (Hb) level determined and classified.

*Corresponding author: E-mail: vickiyanam@gmail.com;

Results: The age of the respondents ranged from 15-49 years with mean and standard deviation of 26.62 ± 6.29 years. Results obtained show that out of 375 pregnant women, 265 had Hb <11g/dl giving 70.67% of anaemia among them. While 18.49% had mild anaemia (Hb:10.0-10.9g/dl), 4.15% had severe anaemia (Hb:<7.0g/dl). Anaemia was statistically associated with rural residence (p=0.024), low educational status (p=0.02), low family income (p=0.003), being married (p=0.016), lower parity (p=0.000), late booking (p=0.001), non-use of family planning (p=0.000), non-use of insecticide-treated nets (p=0.000), febrile illness in index pregnancy (p=0.000) and poor nutrition (p=0.000).

Conclusion: The prevalence of anaemia among pregnant women in the study is high. This has far-reaching negative implications on the health status of the women during pregnancy, delivery and puerperum and that of the fetus. The need for preconception counseling and screening, health and nutrition education, early booking, contraception, treatment and prevention of causes of febrile illness, priority to girl child education and overall poverty eradication measures and recommended.

Keywords: Anaemia; pregnancy; prevalence; southern Nigeria.

1. INTRODUCTION

Anaemia in pregnancy (AIP) is defined by the WHO as haemoglobin concentration of less than 11g/dl (Hb < 11g/dl) in a pregnant women [1]. It is classified into mild, moderate and severe anaemia when the haemoglobin concentration is 10.0 – 10.9g/dl, 7.0-9.9g/dl and < 7.0g/dl respectively [2-6].

Though a preventable morbidity in most cases, AIP has been identified as the commonest pathology affecting pregnant women globally, contributing significantly to maternal, perinatal and infant morbidity and mortality especially in middle and low income countries [1,2,7-9]. Several studies have identified complications associated with AIP including low physical activity, pregnancy-induced hypertension, abortion, high risk of preterm delivery, post-partum haemorrhage, increased need for blood transfusion with the associated risks, impaired immune function, increased cardiac disease, post partum depression and sepsis [4,5,10-14]. Moreover, foetus/infant of anaemic mothers are prone to intrauterine growth retardation, intrauterine death, prematurity, low birth weight (LBW), neural tube defect, low APGAR score at 5 minutes of delivery, developmental delay and impaired physical and cognitive development [5,10-14]. AIP is a contributing factor to obstetric near misses especially in the sub-Saharan African countries; it was also a major cause of the non-attainment of goal 5 of the erstwhile millennium development goals (MDG-5) [6,14-18].

Being the commonest medical morbidity in pregnancy, AIP has a prevalence that cuts across every region of the world with the highest

prevalence found in least developed countries. According to the WHO, as at 2011, about 38.2% of pregnant women globally were anaemic [4,19]. Regionally, sub-Saharan Africa has the highest prevalence of 57%, followed by south east Asia with 48% while south America has the lowest prevalence of 24.1. [14]. The prevalence also varies from country to country and from region to region within same countries. Studies have shown that the prevalence of AIP is as high as 99% among women in tertiary facility in India, [20] 90.5% in urban Pakistan, 97.6% among pregnant adolescents in Ghana [21], 39.9% in south Ethiopia, [8] 47.4% in northern Tanzania [4] to as low as 2.2% among pregnant women with early pregnancy in New Zealand. [22] In Nigeria, the overall prevalence of anaemia among pregnant women is put at 57.8% [1] while studies have documented regional prevalence of 54.5% in Uyo South South Nigeria, [3] 56% in Abakiliki in south east Nigeria, [5] 39.2% in Sokoto north western Nigeria, [7] 76.5% in Abeokuta, south western Nigeria, [12] 71.3% in Azara in north east Nigeria, [23] and 60.4% in a tertiary facility in Ilorin, north central Nigeria. [24]

Anaemia in pregnancy is mostly a deficiency disease caused mainly by iron, folate and vitamin B12 depletion, especially in low income setting. [7,8,25,26] Several risk factors are responsible for the deficiency. These border on medical, economic, social, religious and cultural factors and include parasitic intestinal infestation, malaria, HIV infection and other causes of febrile illness in pregnancy, chronic illnesses, pica, young age, multigravidity, grand multiparity, advanced gestational age, close birth interval, history of excessive blood loss during pregnancy, low socioeconomic status, illiteracy, haemoglobinopathy, low body mass index (BMI) and,

unhealthy cultural dietary practices. [7,8,24,27-31] Almost all the risk factors associated with anaemia in pregnancy are preventable through preconception screening, early booking, use of contraceptives and insecticide-treated nets, deworming, improved nutrition, overall improvement in socioeconomic status, female education and other safe motherhood initiative recommendatios. [13]

This study therefore aimed at determining the prevalence of anaemia at booking as well as identifying and describing risk factors associated with anaemia among pregnant women who presented at a secondary health care facility in a sub-urban area of southern Nigeria.

2. METHODOLOGY

2.1 Study Area

The study was carried out at the antenatal clinic of General Hospital, Etinan, Akwa Ibom State, south-south Nigeria. The General Hospital Etinan was founded as a missionary Hospital in 1927 by Qua Iboe Church (QIC). It was later taken over and is currently run by the Akwa Ibom State Hospital Management Board (HMB). It is a secondary health care facility that takes care of the needs of Etinan community and beyond. It has a maternity unit with a standard antenatal clinic operated by trained nurses supervised by medical officers.

2.2 Study Design

This was a cross-sectional descriptive study involving pregnant women who booked for antenatal care at the maternity unit of the General Hospital, Etinan, Akwa Ibom State, south-south Nigeria.

2.3 Sample Selection

A total of 375 pregnant women who booked at the facility during the period of the study were recruited into the study using the formula: [32]

$$M = \frac{Z^2 p(1-p)}{M^2}$$

where

Z = Confidence level at 95% (standard value of 1.96%), at

M = 5% acceptable margin of error (Standard value 0.05)

P = Prevalence of anaemia among pregnant women in Nigeria (57.8%). [1] Substituting in the above formula gives 375 (in the nearest 10). A non-probability consecutive sampling technique was used to select the respondents.

The inclusion criteria were all consenting pregnant women who booked at the facility during the period of the study. The exclusion criteria were pregnant women with current bleeding, those who came for repeat visit, those with haemoglobinopathies, those who had blood transfusion in the current pregnancy and those who were in labour.

2.4 Data Collection

A pre-tested semi-structured questionnaire was used to collect data from the respondents by the author and trained assistants after due explanation of the purpose of the study, assurance of confidentiality of information and verbal consent obtained from them. Pregnant women with no formal education were assisted by research assistants in the use of local dialect. The data contained in the questionnaire include socio-demographic, family and nutritional characteristics as well as obstetric/medical history of the respondents. A sample of 2mls of blood was collected from the median cubital vein of each pregnant woman into ethylenediaminetetraacetic acid (EDTA) bottle. This was analyzed with haemoglobinometer by trained medical laboratory technician. The haemoglobin level of each respondent was determined and classified according to the WHO criteria into normal haemoglobin (Hb \geq 11.0g/dl); mild anaemia (Hb:10.0-10.9g/dl), moderate anaemia (Hb:7-9.9g/dl) and severe anaemia (Hb < 7.0g/dl).

2.5.1 Data analysis

Data obtained from the study were analysed with Epi-info statistical package (CDC). The percentage of independent and dependent variables were determined. Level of statistical significance was set at $p < 0.05$. Tables and charts were used to display data distribution.

2.6 Ethical Clearance and Consent

Approval for the study was obtained from the Research and Ethical Committee of Akwa Ibom State Ministry of Health. Permission was also obtained from the administrative head of the

hospital and the head of the maternity unit before commencement of data collection. Verbal consent was obtained from the respondents before the administration of the questionnaires and collection of the blood samples.

3. RESULTS

A total of 375 pregnant women were recruited for the study. The results obtained are shown below: Out of 375 pregnant women seen at the facility during the period of the study, majority 265 (70.67%) were anaemic (Hb < 11.0g/dl) while 110(29.33%) had normal haemoglobin level (Hb ≥ 11.0g/dl).

The haemoglobin levels of the anaemic respondents range from 4-10.9g/dl, with mean and standard deviation of 8.7± 1.0g/dl. Fig. 1 depicts the classification of anaemia among the respondents. Out of 265 anaemic respondents, majority, 205 (representing 77.36%) had moderate anaemia (Hb:7-9.9g/dl) while least, 11(representing 4.15%) had severe anaemia (Hb <7.0g/dl).

Table 1 summarises the socio-demographic characteristics of the pregnant women with

anaemia. Their age distribution range from 15-49 years with mean age and standard deviation of 29.62 ± 6.29 years. The majority of the respondents, 80(30.19%) belonged to 30-34 years age bracket while the lowest number, 2(0.75%), belonged to advanced reproductive age (44-49 years). Majority of the respondents, 58.49% came from rural areas. Most of the pregnant women, 159 (59.99%) had lower levels of education (secondary education and below). Also most of the respondents, 163 (61.51%) were married. A higher percentage of the pregnant women were unemployed (56.23%), had lower family income (67.57%) and practised Christianity (92.74%).

Table 2 shows some practices and nutritional characteristics in index pregnancy among the respondents. Majority of the pregnant women did not plan for the pregnancy (58.49%) and did not use insecticide-treated net (ITN), (63.4%) respectively.

Only 20.76% of the pregnant women had good nutritional status during the pregnancy while majority, 79.24%, had poor (fair and poor) nutritional status.




Fig. 1. Classification of anaemia among the respondents

Table 3 shows the obstetric and medical history of the pregnant women in the study. Almost all respondents (97.36%) had lower parity (parity ≤4).

Majority of the pregnant women 233 (87.93%) booked late (Second and third trimesters). A small percentage, 9 (3.4%) of the respondent

had per vaginal bleeding in the index pregnancy. Out of 228 pregnant women who had given birth previously, majority of them 131(57.46%) and 205 (89.91%) had birth interval of 1-2 years and had not used contraceptives respectively. Medically most of the respondent, 88.68% had febrile illness in the index pregnancy while 7.17% of them were HIV positive.

Table 4 shows association between anaemia and socio-demographic characteristics among the pregnant women. Anaemia was statistically associated with rural residence ($p=0.025$), lower educational status ($p = 0.002$), being currently married ($p = 0.016$). and lower monthly family income ($p = 0.0003$).

Table 1. Socio-demographic characteristics of the anaemic respondents

Characteristic	Frequency (N-265)	Percentage (%)
Age (years)		
15-19	15	5.70
20-24	46	17.36
25-29	62	23.40
30-34	80	30.19
35-39	53	20.00
40-44	7	2.64
45-49	2	0.75
Residence:		
Urban	110	41.51
Rural	115	58.49
Level of Education:		
No formal Education	6	2.26
Primary Education	33	12.45
Secondary education	120	45.28
Tertiary education	106	40.00
Currently married:		
Yes	163	61.51
No	102	38.49
Employment status		
Unemployed	116	43.77
Employed	149	56.23
Average Monthly family income (N):		
<50,000	179	67.55
50,000 – 99,000	74	27.92
≥ 100,000	12	4.53
Religion:		
Christianity	259	97.74
Other Religion	6	2.26

Table 2. Some practices and nutritional characteristics of the respondents

Characteristic	Frequency (N-265)	Percentage(%)
Planned for this pregnancy?		
Yes	110	41.51
No	155	58.49
Use of Insecticide-Treated Net (ITN)		
Yes	97	36.60
No	168	63.40
Nutritional Status**		
Good	55	20.76
Fair	200	75.47
Poor	10	3.77

** Nutritional Status:

Good – Eats 3 times a day with good food and meat most days of the week.

Fair – Eats 3 times only occasionally, eats 2 times most days of the week taking meat 2-3 times a week.

Poor – Eats only one to two times most days. Rarely takes meat.

Table 3. Obstetric and medical history of the respondents

Characteristic	Frequency(n=265)	Percentage(%)
Parity:		
0	37	13.96
1-2	170	64.15
3-4	51	19.25
≥ 5	7	2.64
Trimester at booking:		
First (week 1-13)	32	12.07
Second (week -14-28)	141	53.21
Third (week 29-38)	92	34.72
Bleeding per Vaginaan In index pregnancy		
Yes	9	3.4
No	256	96.6
Interval between last and Current pregnancies (n=228)***		
< 6 months	6	2.63
6-11 months	67	29.39
1-2 years	131	>7.46
> 2 years	9	3.95
Uncertain	15	6.58
Use of family planning after Last pregnancy (n=228)		
Yes	23	10.09
No	205	89.91
Febrile illness in index pregnancy (n=265)		
Yes	235	88.68
No	30	11.32
HIV Status (n=265)		
Positive	19	7.17
Negative	246	92.83

***228 of the respondents had parity ≥1

Table 5 summaries the association between AIP and obstetric and medical history and, family and nutritional characteristics of the respondents. AIP was statistically significantly associated with lower parity ($p = 0.000$), late booking ($p = 0.001$), non-use of contraceptives ($p = 0.000$), febrile illness in index pregnancy ($p = 0.000$), poor nutritional status ($p = 0.000$) and non-use of insecticide-treated bed nets ($p = 0.000$)

4. DISCUSSION

The results of the study reveal a high prevalence rate (70.65%) of anaemia among pregnant women seen at General Hospital, Etinan, a sub-urban area of south-south Nigeria. This rate is similar to findings by Idowu, et al., in Abeokuta, western Nigeria, [12] Dattijo, et al., in Azara,

north eastern Nigeria [23] and Huang, et al., in south China. [33] It is however lower than the 99% found by Manjulatha, et al., in a tertiary health facility in India, [20] 90.5% by Baing-Ansari, et al in urban Pakistan, [9] and 76% by Okafor, et al in rural community in Calabar, South-South Nigeria. [34] It higher than findings from other workers in Uyo, south-south, Nigeria, [3] Shagamu, south western Nigeria, [4] Addis – Ababa, Etiopia, [6] south Sudan [25] and Nepal. [35] The high prevalence rate of anaemia obtained in this study with similar and higher rates obtained in other studies are an attestation that anaemia is still a common morbidity among pregnant population across the globe and therefore calls for closer attention and more holistic approach to its prevention, prompt identification and treatment so as to ensure maternal and child well being during and after pregnancy.

Table 4. Association between AIP and socio-demographic characteristics

Characteristic	Anaemia in pregnancy		Statistical Tests and Values
	Yes N(%)	No N(%)	
Age (Year):			
< 30	123(46.45)	45(40.91)	$X^2 = 0.953$
≥ 30	142 (53.59)	65(59.09)	Df 1 p=0.329
Residence:			
Urban	110(41.51)	32(29.09)	$X^2 = 5.095$
Rural	155(58.49)	78(70.91)	Df 1 p = 0.025
Educational level:			
Lower Education	159(59.99)	80(72.73)	$X^2=5.448$
Higher education	106(40.00)	30(27.27)	Df 1 p = 0.020
Currently married			
Yes	163(61.51)	82(79.54)	$X^2 = 5.832$
No	102(38.49)	28(25.46)	Df 1 p = 0.016
Employment status			
Unemployed	116(43.77)	52(47.27)	$X^2 = 0.385$
Employed	149(56.23)	58(52.73)	Df 1 p = 0.535
Religion:			
Christianity	259(97.74)	107(97.27)	$X^2 = 0.071$
Other Religion:	6(2.26)	3(2.73)	Df 1 p = 0.790
Average monthly family Income (in Naira)			
<50,000	179(65.55)	11(82.73)	$X^2 = 0.885$
≥50,000	86(32.45)	19(17.27)	Df 1 p = 0.0003

Although not statistically significant, the study has shown that sociodemographically, pregnant women of higher age (≥ 30 years) were more affected by anaemia than those of lower age (< 30 years). This is in disagreement with findings by Olubukola, et al in a study in Ibadan which found anaemia associated with pregnant women of lower age [36]. The finding here could be due to the fact the population of women above 30 years in the study was more than that of women of lower age. This underscores the need to educate women of advanced reproductive age on practices that could avert anaemia in pregnancy among them.

The study established a significant association between anaemia and other socio-demographic characteristics including rural residence, low educational status, being married and low family monthly income. These findings are similar to findings from several other studies on AIP [8,9,18,37-39]. There is need therefore to embark on a more intensive health education to

women in rural and sub-urban areas, establish more primary health care centres in rural areas to enhance ready access of health workers to rural women for information, education and counseling on measures to prevent anaemia in pregnancy; give priority to girl child education and improve socioeconomic conditions of the rural and sub-urban dwellers.

From Table 5, anaemia was found to be statistically associated with lower parity, late booking, non-use of family planning, febrile illness in index pregnancy, poor nutrition in pregnancy and non-use of insecticide-treated bed nets (ITN). These findings are similar to findings from several other studies on risk factors for anaemia in pregnancy in other settings. [30,31,39-43] This underscores the need to pay special attention to women of lower parity, who sometimes have poor spacing between births, with the aim of educating them on anaemia prevention measures, especially during pregnancy. The need to educate pregnant

women, especially those in the rural areas on the importance of early booking and booking in certified health facilities is also of utmost importance.

Furthermore there is need for early screening of pregnant women for febrile illnesses with the aim

of prompt identification and treatment of causes, especially in sub-Saharan Africa where malaria and other infestations and infections are major causes of fever, haemolysis and resultant anaemia among the general population and pregnant population in particular. [7,27,44-45]

Table 5. Association between AIP and obstetric and medical history and some practices and nutritional characteristics of the respondents

Characteristic	Anaemia in pregnancy		Statistical Tests and Values
	Yes N(%)	No N(%)	
Parity:			
<5	258(97.36)	96(87.27)	$X^2 = 14.958$
≥ 5	7(2.64)	14(12.73)	Df 1 p = 0.000
Trimester at booking:			
<Second Trimester	32(12.07)	29(26.36)	$X^2 = 11.651$
≥Second Trimester	233(89.93)	8(73.64)	Df 1 p = 0.001
Per vaginaam Bleeding In index pregnancy:			
Yes	9(3.4)	7(6.36)	$X^2 = 1.679$
No	256(96.6)	103(93.64)	Df 1 p = 0.195
Interval between last and Index pregnancies (n=228)			
< 2 years	73(32.02)	49(33.33)	$X^2 = 0.070$
≥ 2 years	155(67.99)	98(66.67)	Df 1 p = 0.791
Use of family planning after Last delivery (n=228)			
Yes	23(10.09)	47(31.97)	$X^2 = 28.196$
No	205(89.91)	100(68.03)	Df 1 p = 0.000
Febrile illness in index pregnancy (n=265):			
Yes	235(88.68)	79(71.82)	$X^2 = 16.225$
No	30 (11.32)	31(28.18)	Df 1 p = 0.000
HIV Status (n=265)			
Positive	19(7.17)	7(6.36)	$X^2 = 0.078$
Negative	246(92.83)	103(93.64)	Df 1 p = 0.789
Nutritional Status:			
Good	55(20.76)	49(44.55)	$X^2 = 21.952$
Bad (Fair and Poor)	210(79.24)	61(55.45)	Df 1 p = 0.000
Use of Insecticide-Treated Nets (ITN):			
Yes	97(36.60)	78(70.91)	$X^2 = 36.756$
No	168(63.40)	32(29.09)	Df 1 p = 0.000

Moreover the importance of good nutrition among pregnant women to avert anaemia is a well known fact scientifically. Therefore the need for adequate preconception, conception and post-partum nutrition among women of child bearing age is recommended.

Finally, other risk factors such as non-use of contraceptives and insecticide-treated bed nets (ITN), associated with anaemia in pregnancy identified here, should be appropriately addressed. These cost-effective measures should be made readily available and affordable and possibly free to reproductive aged and pregnant women as important tools to preventing anaemia in pregnancy.

5. LIMITATION

The study concentrated on pregnant women in General Hospital, Etinan only and did not assess pregnant women in the core rural areas of Etinan. Therefore its findings may not be a true reflection on anaemia among pregnant women in Etinan as a whole. Further studies involving core rural areas is recommended.

6. CONCLUSION

The prevalence of anaemia among pregnant women in Etinan, a sub-urban area of south-south Nigeria is high. This has far reaching negative implications for the women during pregnancy, delivery, puerperum and the foetus. The need for preconception counseling, screening, health and nutrition education, early booking, contraception, prompt identification and treatment of causes of febrile illnesses and overall improvement of socio-economic status and priority to girl child education is recommended.

CONSENT

Verbal consent was obtained from all the correspondents.

ETHICAL APPROVAL

All authors hereby declare that the research was approved by the relevant ethical committee and that the research was performed according to ethical standard in compliance with the Declaration of Helsinki.

ACKNOWLEDGEMENT

The authors are grateful to the management and staff of General Hospital, Etinan especially staff

of antenatal clinic for their immeasurable support that resulted in the success of this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Global Health Observation Data Repository/World Health Statistics. Available:<http://aps.who.int/gho/data/node.main.1?ang=en>. Accessed on 12/12/2018.
2. Okube OT, Mirie W, Odhiambo E, Sabira W, Habtu M. Prevalence and factors associated with anaemia among pregnant women attending antenatal clinic in the second and third trimesters at Pumwani Hospital, Kenya. *Open J Obst Gynaecol* 2016;6(1):16-27.
3. Olatunbojun OA, Abasiattai AM, Bassey EA, James RS, Ibanga G, Morgan A. Prevalence of anaemia among pregnant women at booking in the University of Uyo Teaching Hospital, Nigeria. *Biomed Res Int*. Available:<http://doi.org/10.1155/2014/849080>. Accessed on 24/1/2019.
4. Sholeye OO, Animasahun VS, Shoronmu TO. Anaemia in pregnancy and its associated factors among primary care clients in Sagamu, south west Nigeria: A facility-based study. *J Fam Med Prim Care* 2017;6(2):323-329.
5. Ezike ED, Anosie OB, Onoh RC, Nwokpor OS, Umeora OU. The prevalence of anaemia in pregnancy at booking in Abakiliki, Nigeria. *Trop J Obst Gynaecol* 2016;33(3):332-336.
6. Buseri FI, Uko EK, Jeremiah ZA, Usanga EA. Prevalence and risk factors of anaemia among pregnant women in Nigeria. *The Open Haematol J*. 2008;2:14-19.
7. Buhari H, Imoru M, Erhaboro N. Anaemia in pregnant women of Sokoto residents in North Western Nigeria. *J Blood Discord Transfus*. 2016;7:366.
8. Gedefau L, Ayele A, Asre Y, Mosie A. Anaemia and associated factors among pregnant women attending antenatal care clinic in Wolajita, Sodo town, Southern Ethiopia. *Ethiopia J Health*. 2015;25(5): 155-162.

9. Baig Ansar NM, Badruddin SH, Karmaliani R, Harris H, Jehan I, Pusha O, et al. Anaemia prevalence and risk factors in pregnant women in an urban area of Pakistan. *Food Nutr Bulletin*. 2008;29(2): 132-139.
10. Kassa GM, Muche AM, Fekadu GA. Prevalence and determinants of anaemia among pregnant women in Ethiopia: a systemic review and meta-analysis. *BMC Haematol*. 2017;17:17.
11. Nwizu EN, Iliyasu Z, Ibrahim SA, Galadanci HS. Sociodemographic and maternal factors in anaemia in pregnancy at booking in Kano, Northern Nigeria. *Afr J Reprod Health*. 2011;15(4):33-41.
12. Idowu OA, Mafiana CF, Sofiloye D. Anaemia in pregnancy: A survey of pregnant women in Abeokuta Nigeria. *Afr Health Sci*. 2005;5(4):295-299.
13. Islam M. The safe Motherhood initiative and beyond. *Bulletin of the World Health Organization*. Available: <https://www.who.int/> Vol) Accessed on 21/02/19.
14. Stephen G, Mgongo M, Hashim TH, Katanya J, Stray-Pederson B, Msuya SE. Anaemia in pregnancy: Prevalence, risk factors and adverse perinatal outcomes in Northern Tanzania. *Anaemia*. 2018;1-9. 2018/1846280.
15. Oladapo OT, Sule-Odu AO, Olatunji AO, Daniel OJ. "Near Miss" Obstetric events and maternal deaths in Sagamu, Nigeria; a retrospective study. *Reprod Health*. 2005; 2;9.
16. WHO. WHO maternal death and near-miss classifications. Available at: www.who.int. Accessed on 7/2/2019.
17. Tallapureddy S, Velagelati R, Palutia H, Satti CV, "Near Miss" Obstetric events and maternal mortality in a tertiary care hospital. *Indian J Pub Health*. 2017;61(4): 305-308.
18. MDG Monitor. MDG 5: Improve Maternal Health. Available: www.Mdgmonitor.org.
19. Onyeneho NG, Aronu NI, Chukwu N, Agbanodikeizu UP, Chalupowki M, Subramian SV. Factors associated with compliance to recommended micronutrients uptake for prevention of anaemia during pregnancy in urban, periurban and rural communities in south east Nigeria. *J Health Popul Nutr*. 2016; 35:35.
20. Manjulatha B, Padmasri PP, Sravanti TP. Prevalence of anaemia in pregnant women in tertiary care centre. *IOSR J Dental Med Sci*. 2015;14(7):91-95.
21. Intiful FD, Wiredu EK, Asare GA, Asante M, Adjei DN. Anaemia in pregnant adolescent girls with malaria and practicing pica. *Pan Afr Med J*. 2016; 24(96):1-7.
22. Masukume G, Khasan AS, Keny LG, Baker PN, Nelson G. Risk factors and birth outcomes of anaemia in early pregnancy in a nulliparous cohort. *PLOS One* 2015; 10(4):1-15.
23. Dattijo LM, Duru PH, Umar NI. Anaemia in pregnancy: prevalence and associated factors in Azare, North-East Nigeria. *Int J Trop Dis Health*. 2016;11(5):1-9.
24. Babatude AS, Olawunmi HO, Duroloye IA, Shitu AO, Adesina KT, Sani MA. Prevalence of anaemia among pregnant women at antenatal care booking in Ilorin, North Central Nigeria. *Trop J Health Scs* 2017;24(4):1-7.
25. Alemayehu A, Gedefaw L, Yemane T, Asre Y. Prevalence, severity and determinant factors of anaemia among pregnant women in south Sudanese refugees, Pugnido, western Ethiopia. *Anaemia*. 2016; 2016:9817358.
26. Hassan A, Mamman AI, Adaji S, Musa B, Kene S. Anaemia and iron deficiency in pregnant women in Zaria, Nigeria. *Sub-Saharan Afr J Med*. 2014;1:36-39.
27. Anlaakum P, Anto F. Anaemia and associated factors: A cross-sectional study of antenatal attendants at the Sunyani Municipal Hospital, Ghana. *BMC Res Notes*. 2017;10(402):1-8.
28. Kassa GM, Muche AA, Bere AK, Fekadu GA. Prevalence and determinants of anaemia among pregnant women in Ethiopia: A systematic review and meta-analysis. *BMC Haematol*. 2017;17(17):1-11.
29. Lin L, Wei Y, Zhu W, Wang C, Su R, Feng G, et al. Prevalence, risk factors and associated adverse pregnant outcomes of anaemia in Chinese pregnant women: A multicentre retrospective study. *BMC Pregnancy and Child Birth*. 2018;18:111.
30. Alem M, Enawgaw B, Gelaw A, Kena T, Seid M, Olkeba Y. Prevalence of anaemia and associated risk factors among pregnant women attending antenatal care in Azezo Health Centre Gondar town,

- North West Ethiopia. J Interdiscpl Histopath. 2013;1(3):137-144.
31. Lokare PO, Karanjekar VD, Gattani PL, Kulkarni AP. A study of prevalence of anaemia and socio-demographic factors associated with anaemia among pregnant women in Aurangabad City, India. Ann Nig. Med. 2012;6(1):30-34.
 32. Statistics HOW TO. Sample size in statistics. Available: www.statisticshowto.com Accessed on 12-01-2019.
 33. Huang LL, Purvarshi G, Wang SM, Zhong LL, Tang H. The influence of iron-deficiency anaemia during the pregnancy on preterm birth and birth weight in South China. J Food Nutri Res. 2015;3(9):570-574.
 34. Okafor IM, Okpokam DC, Antai AB, Usanga EA. Iron state of pregnant women in rural and urban communities of Cross River State, south-south Nigeria. Niger J Physiol Sci. 2016;31:121-125.
 35. Ghosh S, Trevino JT, Davis D, Shrestha R, Battarai A, Anusree KC, et al. Factors associated with anaemia in pregnant women in Banke, Nepal. FASEB J. 2017; 788:32.
 36. Olubukola A, Odunayo A, Adesina O. Anaemia in pregnancy at two levels of health care in Ibadan, South West Nigeria. Ann Afr Med. 2011;10(4):272-277.
 37. Le CHH. The prevalence of anaemia and moderate – severe anaemia in the US population (NHANES 2003-2012). PLOS ONE. 2016;16665.
 38. Chowdhury HA, Ahmed KR, Jebunessa F, Akter J, Hossain S, Shajaham M. Factors associated with maternal anaemia among pregnant women in Dhata City. BMC Women's Health. 2015;15(77):1-6.
 39. Gwarzo MY, Ugwa EA. The pattern of anaemia among pregnant women in Northern Nigerian. J Med Med Sci. 2013; 4(8):319-323.
 40. Adamgbe M, Balami AD, Bello H. Anaemia and its associated factors among pregnant women in Koko, Kebbi State, Nigeria. Nig J Med. 2017;26(1):29-34.
 41. Tadesse SE, Seid O, Mariam GY, Fekadu A, Wasihun Y, Endris K, et al. Determinants of anaemia among pregnant mothers attending antenatal care in Dessie Town Health facilities, Northern Central Ethiopia: Unmatched case control study. PLOS ONE. 2017;12(3):10-9.
 42. Onoh RC, Lawani OL, Ezeonu PO, Nkwo PO, Onoh TJP, Ajah LO. Predictors of anaemia in pregnancy among pregnant women accessing antenatal care in a poor resource setting in South Eastern Nigeria. Sahel Med J. 2015;18(4):182-187.
 43. Bassi AP, Idoko L, Dibigbo-Ibeaji NM, Ademinyi OG, Ramyil MS, Ogundeko TO, et al. Prevalence of anaemia in pregnancy among women visiting antenatal clinic in Bingham University Teaching Hospital, Jos, Nigeria. Clin Med Res. 2016;5(3):52-62.
 44. Tunkyi K, Moodley J. Anaemia and pregnancy outcomes: A longitudinal study. J Maternal Fetal Neonat Med. 2018; 31(19):2594-2598.
 45. Lamina MA, Sorunmu TO. Prevalence of anaemia in pregnant women attending the antenatal clinic in a Nigerian University Teaching Hospital. Nig Med Pract. 2003; 44(2):39-42.

© 2019 Iyanam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48805>